

Observation of Large CP Violation in the Neutral B Meson System

K. Abe,⁹ K. Abe,³⁷ R. Abe,²⁷ I. Adachi,⁹ Byoung Sup Ahn,¹⁶ H. Aihara,³⁹ M. Akatsu,²⁰ G. Alimonti,⁸ K. Asai,²¹ M. Asai,¹⁰ Y. Asano,⁴⁴ T. Aso,⁴³ V. Aulchenko,² T. Aushev,¹⁴ A. M. Bakich,³⁵ E. Banas,²⁵ S. Behari,⁹ P. K. Behera,⁴⁵ D. Beilene,² A. Bondar,² A. Bozek,²⁵ T. E. Browder,⁸ B. C. K. Casey,⁸ P. Chang,²⁴ Y. Chao,²⁴ K.-F. Chen,²⁴ B. G. Cheon,³⁴ R. Chistov,¹⁴ S.-K. Choi,⁷ Y. Choi,³⁴ L. Y. Dong,¹² J. Dragic,¹⁹ A. Drutskoy,¹⁴ S. Eidelman,² V. Eiges,¹⁴ Y. Enari,²⁰ R. Enomoto,⁹ C. W. Everton,¹⁹ F. Fang,⁸ H. Fujii,⁹ C. Fukunaga,⁴¹ M. Fukushima,¹¹ N. Gabyshev,⁹ A. Garmash,^{2,9} T. J. Gershon,⁹ A. Gordon,¹⁹ K. Gotow,⁴⁶ H. Guler,⁸ R. Guo,²² J. Haba,⁹ H. Hamasaki,⁹ K. Hanagaki,³¹ F. Handa,³⁸ K. Hara,²⁹ T. Hara,²⁹ N. C. Hastings,¹⁹ H. Hayashii,²¹ M. Hazumi,²⁹ E. M. Heenan,¹⁹ Y. Higashino,²⁰ I. Higuchi,³⁸ T. Higuchi,³⁹ T. Hirai,⁴⁰ H. Hirano,⁴² T. Hojo,²⁹ T. Hokuue,²⁰ Y. Hoshi,³⁷ K. Hoshina,⁴² S. R. Hou,²⁴ W.-S. Hou,²⁴ S.-C. Hsu,²⁴ H.-C. Huang,²⁴ Y. Igarashi,⁹ T. Iijima,⁹ H. Ikeda,⁹ K. Ikeda,²¹ K. Inami,²⁰ A. Ishikawa,²⁰ H. Ishino,⁴⁰ R. Itoh,⁹ G. Iwai,²⁷ H. Iwasaki,⁹ Y. Iwasaki,⁹ D. J. Jackson,²⁹ P. Jalocho,²⁵ H. K. Jang,³³ M. Jones,⁸ R. Kagan,¹⁴ H. Kakuno,⁴⁰ J. Kaneko,⁴⁰ J. H. Kang,⁴⁸ J. S. Kang,¹⁶ P. Kapusta,²⁵ N. Katayama,⁹ H. Kawai,³ H. Kawai,³⁹ Y. Kawakami,²⁰ N. Kawamura,¹ T. Kawasaki,²⁷ H. Kichimi,⁹ D. W. Kim,³⁴ Heejong Kim,⁴⁸ H. J. Kim,⁴⁸ Hyunwoo Kim,¹⁶ S. K. Kim,³³ T. H. Kim,⁴⁸ K. Kinoshita,⁵ S. Kobayashi,³² S. Koishi,⁴⁰ H. Konishi,⁴² K. Korotushenko,³¹ P. Krokovny,² R. Kulasiri,⁵ S. Kumar,³⁰ T. Kuniya,³² E. Kurihara,³ A. Kuzmin,² Y.-J. Kwon,⁴⁸ J. S. Lange,⁶ G. Leder,¹³ M. H. Lee,⁹ S. H. Lee,³³ C. Leonidopoulos,³¹ Y.-S. Lin,²⁴ D. Liventsev,¹⁴ R.-S. Lu,²⁴ J. MacNaughton,¹³ D. Marlow,³¹ T. Matsubara,³⁹ S. Matsui,²⁰ S. Matsumoto,⁴ T. Matsumoto,²⁰ Y. Mikami,³⁸ K. Misono,²⁰ K. Miyabayashi,²¹ H. Miyake,²⁹ H. Miyata,²⁷ L. C. Moffitt,¹⁹ G. R. Moloney,¹⁹ G. F. Moorhead,¹⁹ S. Mori,⁴⁴ T. Mori,⁴ A. Murakami,³² T. Nagamine,³⁸ Y. Nagasaka,¹⁰ Y. Nagashima,²⁹ T. Nakadaira,³⁹ T. Nakamura,⁴⁰ E. Nakano,²⁸ M. Nakao,⁹ H. Nakazawa,⁴ J. W. Nam,³⁴ Z. Natkaniec,²⁵ K. Neichi,³⁷ S. Nishida,¹⁷ O. Nitoh,⁴² S. Noguchi,²¹ T. Nozaki,⁹ S. Ogawa,³⁶ T. Ohshima,²⁰ Y. Ohshima,⁴⁰ T. Okabe,²⁰ T. Okazaki,²¹ S. Okuno,¹⁵ S. L. Olsen,⁸ H. Ozaki,⁹ P. Pakhlov,¹⁴ H. Palka,²⁵ C. S. Park,³³ C. W. Park,¹⁶ H. Park,¹⁸ L. S. Peak,³⁵ M. Peters,⁸ L. E. Piilonen,⁴⁶ E. Prebys,³¹ J. L. Rodriguez,⁸ N. Root,² M. Rozanska,²⁵ K. Rybicki,²⁵ J. Ryuko,²⁹ H. Sagawa,⁹ Y. Sakai,⁹ H. Sakamoto,¹⁷ M. Satapathy,⁴⁵ A. Satpathy,^{9,5} S. Schrenk,⁵ S. Semenov,¹⁴ K. Senyo,²⁰ Y. Settai,⁴ M. E. Sevior,¹⁹ H. Shibuya,³⁶ B. Shwartz,² A. Sidorov,² S. Stanić,⁴⁴ A. Sugi,²⁰ A. Sugiyama,²⁰ K. Sumisawa,⁹ T. Sumiyoshi,⁹ J.-I. Suzuki,⁹ K. Suzuki,³ S. Suzuki,⁴⁷ S. Y. Suzuki,⁹ S. K. Swain,⁸ H. Tajima,³⁹ T. Takahashi,²⁸ F. Takasaki,⁹ M. Takita,²⁹ K. Tamai,⁹ N. Tamura,²⁷ J. Tanaka,³⁹ M. Tanaka,⁹ G. N. Taylor,¹⁹ Y. Teramoto,²⁸ M. Tomoto,⁹ T. Tomura,³⁹ S. N. Tovey,¹⁹ K. Trabelsi,⁸ T. Tsuboyama,⁹ T. Tsukamoto,⁹ S. Uehara,⁹ K. Ueno,²⁴ Y. Unno,³ S. Uno,⁹ Y. Ushiroda,⁹ S. E. Vahsen,³¹ K. E. Varvell,³⁵ C. C. Wang,²⁴ C. H. Wang,²³ J. G. Wang,⁴⁶ M.-Z. Wang,²⁴ Y. Watanabe,⁴⁰ E. Won,³³ B. D. Yabsley,⁹ Y. Yamada,⁹ M. Yamaga,³⁸ A. Yamaguchi,³⁸ H. Yamamoto,⁸ T. Yamanaka,²⁹ Y. Yamashita,²⁶ M. Yamauchi,⁹ S. Yanaka,⁴⁰ J. Yashima,⁹ M. Yokoyama,³⁹ K. Yoshida,²⁰ Y. Yusa,³⁸ H. Yuta,¹ C. C. Zhang,¹² J. Zhang,⁴⁴ H. W. Zhao,⁹ Y. Zheng,⁸ V. Zhilich,² and D. Žontar⁴⁴

(Belle Collaboration)

¹Aomori University, Aomori

²Budker Institute of Nuclear Physics, Novosibirsk

³Chiba University, Chiba

⁴Chuo University, Tokyo

⁵University of Cincinnati, Cincinnati, Ohio

⁶University of Frankfurt, Frankfurt

⁷Gyeongsang National University, Chinju

⁸University of Hawaii, Honolulu, Hawaii

⁹High Energy Accelerator Research Organization (KEK), Tsukuba

¹⁰Hiroshima Institute of Technology, Hiroshima

¹¹Institute for Cosmic Ray Research, University of Tokyo, Tokyo

¹²Institute of High Energy Physics, Chinese Academy of Sciences, Beijing

¹³Institute of High Energy Physics, Vienna

¹⁴Institute for Theoretical and Experimental Physics, Moscow

¹⁵Kanagawa University, Yokohama

¹⁶Korea University, Seoul

¹⁷Kyoto University, Kyoto

¹⁸Kyungpook National University, Taegu

¹⁹University of Melbourne, Victoria

- ²⁰Nagoya University, Nagoya
²¹Nara Women's University, Nara
²²National Kaohsiung Normal University, Kaohsiung
²³National Lien-Ho Institute of Technology, Miao Li
²⁴National Taiwan University, Taipei
²⁵H. Niewodniczanski Institute of Nuclear Physics, Krakow
²⁶Nihon Dental College, Niigata
²⁷Niigata University, Niigata
²⁸Osaka City University, Osaka
²⁹Osaka University, Osaka
³⁰Panjab University, Chandigarh
³¹Princeton University, Princeton, New Jersey
³²Saga University, Saga
³³Seoul National University, Seoul
³⁴Sungkyunkwan University, Suwon
³⁵University of Sydney, Sydney, New South Wales
³⁶Toho University, Funabashi
³⁷Tohoku Gakuin University, Tagajo
³⁸Tohoku University, Sendai
³⁹University of Tokyo, Tokyo
⁴⁰Tokyo Institute of Technology, Tokyo
⁴¹Tokyo Metropolitan University, Tokyo
⁴²Tokyo University of Agriculture and Technology, Tokyo
⁴³Toyama National College of Maritime Technology, Toyama
⁴⁴University of Tsukuba, Tsukuba
⁴⁵Utkal University, Bhubaneswar
⁴⁶Virginia Polytechnic Institute and State University, Blacksburg, Virginia
⁴⁷Yokkaichi University, Yokkaichi
⁴⁸Yonsei University, Seoul

(Received 18 July 2001; published 14 August 2001)

We present a measurement of the standard model CP violation parameter $\sin 2\phi_1$ based on a 29.1 fb^{-1} data sample collected at the $Y(4S)$ resonance with the Belle detector at the KEKB asymmetric-energy e^+e^- collider. One neutral B meson is fully reconstructed as a $J/\psi K_S$, $\psi(2S)K_S$, $\chi_{c1}K_S$, $\eta_c K_S$, $J/\psi K_L$, or $J/\psi K^{*0}$ decay and the flavor of the accompanying B meson is identified from its decay products. From the asymmetry in the distribution of the time intervals between the two B meson decay points, we determine $\sin 2\phi_1 = 0.99 \pm 0.14(\text{stat}) \pm 0.06(\text{syst})$. We conclude that we have observed CP violation in the neutral B meson system.

DOI: 10.1103/PhysRevLett.87.091802

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Kobayashi and Maskawa (KM) proposed, in 1973, a model where CP violation is incorporated as an irreducible complex phase in the weak-interaction quark mixing matrix [1]. The idea, which was presented at a time when only the u , d , and s quarks were known to exist, was remarkable because it required the existence of six quarks. The subsequent discoveries of the c , b , and t quarks, and the compatibility of the model with the CP violation observed in the neutral K meson system led to the incorporation of the KM mechanism into the standard model, even though it had not been conclusively tested experimentally.

In 1981, Sanda, Bigi, and Carter [2] pointed out that the KM model predicted large CP violation in certain decays of B mesons for a range of quark mixing parameters. Subsequent measurements of the B meson lifetime [3] and the discovery of $B^0\bar{B}^0$ mixing [4] indicated that the parameters lie within such a range. Thus, measurements of CP violation in B meson decays provide important tests of the KM model.

The model predicts a CP violating asymmetry in the time-dependent rates for initial B^0 and \bar{B}^0 decays to a common CP eigenstate, f_{CP} [2]. In the case where $f_{CP} = (c\bar{c})K^0$, the asymmetry is given by

$$A(t) \equiv \frac{\Gamma(\bar{B}^0 \rightarrow f_{CP}) - \Gamma(B^0 \rightarrow f_{CP})}{\Gamma(\bar{B}^0 \rightarrow f_{CP}) + \Gamma(B^0 \rightarrow f_{CP})} \\ = -\xi_f \sin 2\phi_1 \sin \Delta m_d t,$$

where $\Gamma[\bar{B}^0(B^0) \rightarrow f_{CP}]$ is the decay rate for $\bar{B}^0(B^0)$ to f_{CP} at a proper time t after production, ξ_f is the CP eigenvalue of f_{CP} , Δm_d is the mass difference between the two B^0 mass eigenstates, and ϕ_1 is one of the three internal angles of the unitarity triangle, defined as $\phi_1 \equiv \pi - \arg\left(\frac{-V_{tb}^* V_{td}}{V_{cb}^* V_{cd}}\right)$ [5]. For the $(c\bar{c})K^0$ decays, both the ambiguity due to strong interactions and the contribution from direct CP violation are expected to be small [5].

Our previous determination, using a data sample taken in 1999–2000, found $\sin 2\phi_1 = 0.58^{+0.32}_{-0.34}(\text{stat})^{+0.09}_{-0.10}(\text{syst})$

[6], which is consistent with the KM model constraints from indirect measurements [7]. Although the combination of this result with other measurements of $\sin 2\phi_1$ [8] strongly indicates violation of CP symmetry in B meson decays, the published results are still not conclusive. In this Letter we report a new measurement of $\sin 2\phi_1$ that uses improved reconstruction algorithms and incorporates data taken in 2001 to achieve a fourfold increase in the size of the event sample. The result reported here includes the earlier data and supersedes the previous value. All data samples have been analyzed and reconstructed with the same consistent procedure.

We use a 29.1 fb^{-1} data sample, which contains $31.3 \times 10^6 \overline{B}B$ pairs, collected with the Belle detector at the KEKB asymmetric-energy e^+e^- (3.5 on 8 GeV) collider [9]. KEKB operates at the $Y(4S)$ resonance ($\sqrt{s} = 10.58 \text{ GeV}$) with a peak luminosity that exceeds $4 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$. The Belle detector is a large-solid-angle magnetic spectrometer that consists of a three-layer silicon vertex detector (SVD), a 50-layer central drift chamber (CDC), a mosaic of aerogel threshold Čerenkov counters (ACC), time-of-flight scintillation counters (TOF), and an array of CsI(Tl) crystals (ECL) located inside a superconducting solenoid coil that provides a 1.5 T magnetic field. An iron flux return located outside of the coil is instrumented to detect K_L mesons and to identify muons (KLM). The detector is described in detail elsewhere [10].

We measure $\sin 2\phi_1$ using $B^0\overline{B}^0$ meson pairs produced at the $Y(4S)$ resonance. The two mesons remain in a coherent p -wave state until one of them decays. The decay of one of the B mesons at time t_{tag} to a final state, f_{tag} , which distinguishes between B^0 and \overline{B}^0 , projects the accompanying B meson onto the opposite b flavor at t_{tag} ; this meson decays to f_{CP} at time t_{CP} . CP violation manifests itself as an asymmetry $A(\Delta t)$, where Δt is the proper time interval $\Delta t \equiv t_{CP} - t_{\text{tag}}$. At KEKB, the $Y(4S)$ is produced with a Lorentz boost of $\beta\gamma = 0.425$ nearly along the electron beam line (z). Since the B^0 and \overline{B}^0 mesons are nearly at rest in the $Y(4S)$ center of mass system (cms), Δt can be determined from the displacement in z between the f_{CP} and f_{tag} decay vertices—i.e., $\Delta t \simeq (z_{CP} - z_{\text{tag}})/\beta\gamma c \equiv \Delta z/\beta\gamma c$.

The measurement requires the reconstruction of $B^0 \rightarrow f_{CP}$ decays, the determination of the b flavor of the accompanying (tagging) B meson, the measurement of Δt , and a fit of the expected Δt distribution to the measured distribution using a likelihood method.

We reconstruct B^0 decays to the following CP eigenstates [11]: $J/\psi K_S$, $\psi(2S)K_S$, $\chi_{c1}K_S$, $\eta_c K_S$ for $\xi_f = -1$ and $J/\psi K_L$ for $\xi_f = +1$. We also use $B^0 \rightarrow J/\psi K^{*0}$ decays where $K^{*0} \rightarrow K_S\pi^0$. Here the final state is a mixture of even and odd CP , depending on the relative orbital angular momentum of the J/ψ and K^{*0} . The CP content is determined from a fit to the full angular distribution of all $J/\psi K^*$ decay modes other than $K^{*0} \rightarrow K_S\pi^0$. We find that the final state is primarily $\xi_f = +1$; the $\xi_f = -1$ fraction is $0.19 \pm 0.04(\text{stat}) \pm 0.04(\text{syst})$ [12].

J/ψ and $\psi(2S)$ mesons are reconstructed via their decays to $\ell^+\ell^-$ ($\ell = \mu, e$). The $\psi(2S)$ is also reconstructed via $J/\psi\pi^+\pi^-$, and the χ_{c1} via $J/\psi\gamma$. The η_c is detected in the $K^+K^-\pi^0$ and $K_S K^-\pi^+$ modes. For the $J/\psi K_S$ mode, we use $K_S \rightarrow \pi^+\pi^-$ and $\pi^0\pi^0$ decays; for other modes we use only $K_S \rightarrow \pi^+\pi^-$.

The J/ψ , $\psi(2S)$, and K_S selection has been described elsewhere [6]. For $\chi_{c1}K_S$ decays, we select $\chi_{c1} \rightarrow J/\psi\gamma$ decays, rejecting γ 's that are consistent with $\pi^0 \rightarrow \gamma\gamma$ decays, and use the requirement $385 < M_{\gamma\ell\ell} - M_{\ell\ell} < 430.5 \text{ MeV}/c^2$. For η_c decays, we distinguish kaons from pions using a combination of CDC energy loss measurements, flight times measured in the TOF, and the response of the ACC. Candidate $\eta_c \rightarrow K^+K^-\pi^0(K_S K^-\pi^+)$ decays are selected with a $KK\pi$ mass requirement that takes into account the natural width of the η_c . For $J/\psi K^{*0}(K_S\pi^0)$ decays, we use $K_S\pi^0$ combinations that have an invariant mass within $75 \text{ MeV}/c^2$ of the nominal K^* mass. We reduce background from low-momentum π^0 's by requiring $\cos\theta_{K^*} < 0.8$, where θ_{K^*} is the angle between the K_S momentum vector and the K^{*0} flight direction calculated in the K^{*0} rest frame.

We identify B decays using the energy difference $\Delta E \equiv E_B^{\text{cms}} - E_{\text{beam}}^{\text{cms}}$ and the beam-energy constrained mass $M_{bc} \equiv \sqrt{(E_{\text{beam}}^{\text{cms}})^2 - (p_B^{\text{cms}})^2}$, where $E_{\text{beam}}^{\text{cms}}$ is the cms beam energy, and E_B^{cms} and p_B^{cms} are the cms energy and momentum of the B candidate.

Figure 1 shows the combined M_{bc} distribution for all channels other than $J/\psi K_L$ after a mode-dependent requirement on ΔE . The B meson signal region is defined as $5.270 < M_{bc} < 5.290 \text{ GeV}/c^2$. Table I lists the numbers of observed candidates (N_{ev}) and the background (N_{bkgd}) determined by extrapolating the rate in the nonsignal ΔE vs M_{bc} region into the signal region.

Candidate $B^0 \rightarrow J/\psi K_L$ decays are selected by requiring ECL and/or KLM hit patterns that are consistent with the presence of a shower induced by a neutral hadron. The centroid of the shower is required to be in a 45° cone centered on the K_L direction that is inferred from two-body decay kinematics and the measured four-momentum of the J/ψ . We reduce the background by means of a likelihood ratio that depends on the J/ψ cms momentum, the angle




FIG. 1. The beam-energy constrained mass distribution for all decay modes combined other than $J/\psi K_L$. The shaded area is the estimated background. The signal region is the range $5.27\text{--}5.29 \text{ GeV}/c^2$.

TABLE I. The numbers of observed events (N_{ev}) and the estimated background (N_{bkgd}) in the signal region for each f_{CP} mode.

Mode	N_{ev}	N_{bkgd}
$J/\psi(\ell^+\ell^-)K_S(\pi^+\pi^-)$	457	11.9
$J/\psi(\ell^+\ell^-)K_S(\pi^0\pi^0)$	76	9.4
$\psi(2S)(\ell^+\ell^-)K_S(\pi^+\pi^-)$	39	1.2
$\psi(2S)(J/\psi\pi^+\pi^-)K_S(\pi^+\pi^-)$	46	2.1
$\chi_{c1}(J/\psi\gamma)K_S(\pi^+\pi^-)$	24	2.4
$\eta_c(K^+K^-\pi^0)K_S(\pi^+\pi^-)$	23	11.3
$\eta_c(K_S K^-\pi^+)K_S(\pi^+\pi^-)$	41	13.6
$J/\psi K^{*0}(K_S\pi^0)$	41	6.7
Subtotal	747	58.6
$J/\psi(\ell^+\ell^-)K_L$	569	223

between the K_L and its nearest-neighbor charged track, the charged track multiplicity of the event, the extent to which the event is consistent with a $B^+ \rightarrow J/\psi K^{*+}(K_L\pi^+)$ hypothesis, and the polar angle with respect to the z direction of the reconstructed B^0 meson in the cms. In addition, we remove events that are reconstructed as $B^0 \rightarrow J/\psi K_S$, $J/\psi K^{*0}(K^+\pi^-, K_S\pi^0)$, $B^+ \rightarrow J/\psi K^+$, or $J/\psi K^{*+}(K^+\pi^0, K_S\pi^+)$ decays. Finally, K_L clusters with positions that match photons from reconstructed π^0 's are also rejected.

Figure 2 shows the p_B^{cms} distribution, calculated with the $B^0 \rightarrow J/\psi K_L$ two-body decay hypothesis. The histograms are the results of a fit to the signal and background distributions. The shapes are derived from Monte Carlo (MC) simulations [13], and the normalization and peak position of the signal are allowed to vary. There are 397 entries in the $0.2 \leq p_B^{cms} \leq 0.45$ GeV/ c signal region with KLM clusters. There are 172 entries in the range $0.2 \leq p_B^{cms} \leq 0.40$ GeV/ c with clusters in the ECL only. The fit finds a total of $(346 \pm 29)J/\psi K_L$ signal events, and a signal purity of 61%.

Leptons, charged pions, and kaons that are not associated with a reconstructed CP eigenstate decay are used to identify the flavor of the accompanying B meson. Initially, the b -flavor determination is performed at the track level.




FIG. 2. The p_B^{cms} distribution for $B^0 \rightarrow J/\psi K_L$ candidates with the results of the fit. The solid line is the signal plus background; the shaded area is background only. The signal region for KLM (ECL-only) clusters is $0.2 \leq p_B^{cms} \leq 0.45(0.40)$ GeV/ c .

Several categories of well measured tracks that distinguish the b flavor by the track's charge are selected: high momentum leptons from $b \rightarrow c\ell^-\bar{\nu}$, lower momentum leptons from $c \rightarrow s\ell^+\nu$, charged kaons and Λ baryons from $b \rightarrow c \rightarrow s$, high momentum pions that originate from decays of the type $B^0 \rightarrow D^{(*)-}(\pi^+, \rho^+, a_1^+, \text{etc.})$, and slow pions from $D^{*-} \rightarrow \bar{D}^0\pi^-$. We use the MC to determine a category-dependent variable that indicates whether a track originates from a B^0 or \bar{B}^0 . The values of this variable range from -1 for a reliably identified \bar{B}^0 to $+1$ for a reliably identified B^0 and depend on the tagging particle's charge, cms momentum, polar angle, and particle-identification probability, as well as other kinematic and event shape quantities. The results from the separate track categories are then combined to take into account correlations in the case of multiple track-level tags. This stage determines two event-level parameters, q and r . The first, q , has the discrete values $q = +1$ when the tag-side B meson is more likely to be a B^0 and -1 when it is more likely to be a \bar{B}^0 . The parameter r is an event-by-event flavor-tagging dilution factor which ranges from $r = 0$ for no flavor discrimination to $r = 1$ for unambiguous flavor assignment. It is used only to sort data into six intervals of r , according to flavor purity; the wrong-tag probabilities for the final fit are determined from data.

The probabilities of an incorrect flavor assignment, w_l ($l = 1, 6$), are determined directly from the data for the six r intervals using exclusively reconstructed, self-tagged $B^0 \rightarrow D^{*-}\ell^+\nu$, $D^{(*)-}\pi^+$, $D^{*-}\rho^+$, and $J/\psi K^{*0}(K^+\pi^-)$ decays. The b flavor of the accompanying B meson is assigned according to the flavor-tagging algorithm described above. The exclusive decay and tag vertices are reconstructed using the same vertexing algorithm that is used in the analysis to measure CP asymmetry. The values of w_l are obtained from the amplitudes of the time-dependent $B^0\bar{B}^0$ mixing oscillations: $(N_{OF} - N_{SF})/(N_{OF} + N_{SF}) = (1 - 2w_l)\cos(\Delta m_d\Delta t)$. Here N_{OF} and N_{SF} are the numbers of opposite and same flavor events. We fix Δm_d at the world average value [14]. Table II lists the resulting w_l values together with the fraction of the events (f_l) in each r interval. The total effective tagging efficiency is $\sum_l f_l(1 - 2w_l)^2 = 0.270 \pm 0.008(\text{stat})_{-0.009}^{+0.006}(\text{syst})$.

The vertex positions for the f_{CP} and f_{tag} decays are reconstructed using tracks that have at least one

TABLE II. The event fractions (f_l) and incorrect flavor assignment probabilities (w_l) for each r interval. The errors include both statistical and systematic uncertainties.

l	r	f_l	w_l
1	0.000–0.250	0.405	$0.465_{-0.009}^{+0.010}$
2	0.250–0.500	0.149	$0.352_{-0.014}^{+0.015}$
3	0.500–0.625	0.081	$0.243_{-0.030}^{+0.021}$
4	0.625–0.750	0.099	$0.176_{-0.017}^{+0.022}$
5	0.750–0.875	0.123	$0.110_{-0.014}^{+0.022}$
6	0.875–1.000	0.140	$0.041_{-0.010}^{+0.011}$

three-dimensional coordinate determined from associated r - ϕ and z hits in the same SVD layer along with one or more additional z hits in the other layers. Each vertex position is required to be consistent with the interaction point profile smeared in the r - ϕ plane by the B meson decay length. The f_{CP} vertex is determined using lepton tracks from J/ψ or $\psi(2S)$ decays, or prompt tracks from η_c decays. The f_{tag} vertex is determined from well reconstructed tracks not assigned to f_{CP} . Tracks that form a K_S are not used. The MC indicates that the typical vertex-finding efficiency and vertex resolution (rms) for z_{CP} (z_{tag}) are 92 (91)% and 75 (140) μm , respectively.

The proper-time interval resolution for the signal, $R_{sig}(\Delta t)$, is obtained by convolving a sum of two Gaussians (a *main* component due to the SVD vertex resolution and charmed meson lifetimes, plus a *tail* component caused by poorly reconstructed tracks) with a function that takes into account the cms motion of the B mesons. The fraction in the main Gaussian is determined to be 0.97 ± 0.02 from a study of $B^0 \rightarrow D^{*-}\pi^+$, $D^{*-}\rho^+$, $D^-\pi^+$, $J/\psi K^{*0}$, $J/\psi K_S$, and $B^+ \rightarrow \bar{D}^0\pi^+$, $J/\psi K^+$ events. The means (μ_{main} , μ_{tail}) and widths (σ_{main} , σ_{tail}) of the Gaussians are calculated event-by-event from the f_{CP} and f_{tag} vertex fit error matrices and the χ^2 values of the fit; typical values are $\mu_{main} = -0.24$ ps, $\mu_{tail} = 0.18$ ps and $\sigma_{main} = 1.49$ ps, $\sigma_{tail} = 3.85$ ps. The background resolution $R_{bkg}(\Delta t)$ has the same functional form but the parameters are obtained from a sideband region in M_{bc} and ΔE . We obtain lifetimes for the neutral and charged B mesons using the same procedure; the results [15] agree well with the world average values.

After vertexing we find 560 events with $q = +1$ flavor tags and 577 events with $q = -1$. Figure 3 shows the observed Δt distributions for the $q\xi_f = +1$ (solid points) and $q\xi_f = -1$ (open points) event samples. There is a clear asymmetry between the two distributions; this demonstrates that CP symmetry is violated.

We determine $\sin 2\phi_1$ by performing an unbinned maximum-likelihood fit of a CP violating probability




FIG. 3. Δt distributions for the events with $q\xi_f = +1$ (solid points) and $q\xi_f = -1$ (open points). The results of the global fit (with $\sin 2\phi_1 = 0.99$) are shown as solid and dashed curves, respectively.

density function (pdf) to the observed Δt distributions. For modes other than $J/\psi K^{*0}$ the pdf expected for the signal is

$$\mathcal{P}_{sig}(\Delta t, q, w_l, \xi_f) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{2\tau_{B^0}} \{1 - \xi_f q(1 - 2w_l) \times \sin 2\phi_1 \sin(\Delta m_d \Delta t)\},$$

where we fix τ_{B^0} and Δm_d at their world average values [14]. The pdf used for the background distribution is $\mathcal{P}_{bkg}(\Delta t) = f_\tau e^{-|\Delta t|/\tau_{bkg}}/2\tau_{bkg} + (1 - f_\tau)\delta(\Delta t)$, where f_τ is the fraction of the background component with an effective lifetime τ_{bkg} and δ is the Dirac delta function. For all f_{CP} modes other than $J/\psi K_L$, a study using events in background-dominated regions of ΔE vs M_{bc} shows that f_τ is negligibly small. For these modes, $\mathcal{P}_{bkg}(\Delta t) = \delta(\Delta t)$.

The $J/\psi K_L$ background is dominated by $B \rightarrow J/\psi X$ decays where some final states are CP eigenstates. We estimate the fractions of the background components with and without a true K_L cluster by fitting the p_B^{cms} distribution to the expected shapes determined from the MC. We also use the MC to determine the fraction of events with definite CP content within each component.

The result is a background that is 71% non- CP modes with $\tau_{bkg} = \tau_B$. For the CP -mode backgrounds we use the signal pdf given above with the appropriate ξ_f values. For $J/\psi K^*(K_L\pi^0)$, which is 13% of the background, we use the $\xi_f = -1$ content determined from the full $J/\psi K^*$ sample. The remaining backgrounds are $\xi_f = -1$ states (10%) including $J/\psi K_S$, and $\xi_f = +1$ states (5%) including $\psi(2S)K_L$, $\chi_{c1}K_L$, and $J/\psi\pi^0$.

For the $J/\psi K^*$ mode, we include the Δt and transversity angle θ_{tr} [16] distributions in the likelihood [12]. We use the ξ_f content determined from the full angular analysis.

Each pdf is convolved with the appropriate $R(\Delta t)$ to determine the likelihood value for each event as a function of $\sin 2\phi_1$:

$$P_i = \int \{f_{sig} \mathcal{P}_{sig}(\Delta t', q, w_l, \xi_f) R_{sig}(\Delta t - \Delta t') + (1 - f_{sig}) \mathcal{P}_{bkg}(\Delta t') R_{bkg}(\Delta t - \Delta t')\} d\Delta t',$$

where f_{sig} is the probability that the event is signal, calculated as a function of p_B^{cms} for $J/\psi K_L$ and of ΔE and M_{bc} for other modes. The only free parameter is $\sin 2\phi_1$, which is determined by maximizing the likelihood function $L = \prod_i P_i$, where the product is over all events.

The result of the fit is

$$\sin 2\phi_1 = 0.99 \pm 0.14(\text{stat}) \pm 0.06(\text{sys}).$$

In Fig. 4(a) we show the asymmetries for the combined data sample that are obtained by applying the fit to the events in each Δt bin separately. The smooth curve is the result of the global unbinned fit. Figures 4(b) and 4(c) show the corresponding asymmetries for the $(c\bar{c})K_S$ ($\xi_f = -1$) and the $J/\psi K_L$ ($\xi_f = +1$) modes separately. The observed asymmetries for the different CP states are opposite, as expected. The curves are the results of unbinned




FIG. 4. (a) The asymmetry obtained from separate fits to each Δt bin for the full data sample; the curve is the result of the global fit. The corresponding plots for the (b) $(c\bar{c})K_S$ ($\xi_f = -1$), (c) $J/\psi K_L$ ($\xi_f = +1$), and (d) B^0 control samples are also shown. The curves are the results of the fit applied separately to the individual data samples.

fits applied separately to the two samples; the resultant $\sin 2\phi_1$ values are $0.84 \pm 0.17(\text{stat})$ and $1.31 \pm 0.23(\text{stat})$, respectively.

The systematic error is dominated by uncertainties due to effects of the tails of the vertex distributions, which contribute 0.04. Other significant contributions come from uncertainties (a) in w_l (0.03); (b) in the resolution function parameters (0.02); and (c) in the $J/\psi K_L$ background fraction (0.02). The errors introduced by uncertainties in Δm_d and τ_{B^0} are 0.01 or less.

We performed a number of checks on the measurement. Table III lists the results obtained by applying the same analysis to various subsamples. All values are statistically consistent with each other. The result is unchanged if we use the w_l 's determined separately for $f_{\text{tag}} = B^0$ and \bar{B}^0 . Fitting to the non-CP eigenstate self-tagged modes $B^0 \rightarrow D^{(*)-} \pi^+$, $D^{*-} \rho^+$, $J/\psi K^{*0}(K^+ \pi^-)$, and $D^{*-} \ell^+ \nu$, where no asymmetry is expected, yields 0.05 ± 0.04 . The asymmetry distribution for this control sample is shown in Fig. 4(d). As a further check, we used three independent

TABLE III. The values of $\sin 2\phi_1$ for various subsamples (statistical errors only).

Sample	$\sin 2\phi_1$
$f_{\text{tag}} = B^0 (q = +1)$	0.84 ± 0.21
$f_{\text{tag}} = \bar{B}^0 (q = -1)$	1.11 ± 0.17
$J/\psi K_S (\pi^+ \pi^-)$	0.81 ± 0.20
$(c\bar{c})K_S$ except $J/\psi K_S (\pi^+ \pi^-)$	1.00 ± 0.40
$J/\psi K_L$	1.31 ± 0.23
$J/\psi K^{*0} (K_S \pi^0)$	0.85 ± 1.45
All	0.99 ± 0.14

CP fitting programs and two different algorithms for the f_{tag} vertexing and found no discrepancy.

We conclude that there is large CP violation in the neutral B meson system. A zero value for $\sin 2\phi_1$ is ruled out at a level greater than 6σ . Our result is consistent with the higher range of values allowed by the constraints of the KM model as well as with our previous measurement.

We thank the KEKB accelerator group for the excellent operation of the KEKB accelerator. We acknowledge support from the Ministry of Education, Culture, Sports, Science and Technology of Japan and the Japan Society for the Promotion of Science; the Australian Research Council and the Australian Department of Industry, Science and Resources; the Department of Science and Technology of India; the BK21 program of the Ministry of Education of Korea and the Center for High Energy Physics sponsored by the KOSEF; the Polish State Committee for Scientific Research under Contract No. 2P03B 17017; the Ministry of Science and Technology of Russian Federation; the National Science Council and the Ministry of Education of Taiwan; and the U.S. Department of Energy.

Note added.—As we were preparing to submit this paper, we became aware of a paper from the BABAR Collaboration [17] which also reports on CP violation in the B meson system.

- [1] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [2] A. B. Carter and A. I. Sanda, Phys. Rev. D **23**, 1567 (1981); I. I. Bigi and A. I. Sanda, Nucl. Phys. **B193**, 85 (1981).
- [3] MAC Collaboration, E. Fernandez *et al.*, Phys. Rev. Lett. **51**, 1022 (1983); Mark II Collaboration, N. Lockyer *et al.*, Phys. Rev. Lett. **51**, 1316 (1983).
- [4] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **192**, 245 (1987).
- [5] H. Quinn and A. I. Sanda, Eur. Phys. J. C **15**, 626 (2000). (This angle is also known as β .)
- [6] Belle Collaboration, A. Abashian *et al.*, Phys. Rev. Lett. **86**, 2509 (2001).
- [7] For example, M. Ciuchini *et al.*, hep-ph/0012308.
- [8] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **86**, 2515 (2001), reports $\sin 2\phi_1 = 0.34 \pm 0.20 \pm 0.05$; CDF Collaboration, T. Affolder *et al.*, Phys. Rev. D **61**, 072005 (2000), reports $\sin 2\phi_1 = 0.79^{+0.41}_{-0.44}$; ALEPH Collaboration, R. Barate *et al.*, Phys. Lett. B **492**, 259 (2000),

- reports $\sin 2\phi_1 = 0.84_{-1.04}^{+0.82} \pm 0.16$; OPAL Collaboration, K. Ackerstaff *et al.*, Eur. Phys. J. C **5**, 379 (1998), reports $\sin 2\phi_1 = 3.2_{-2.0}^{+1.8} \pm 0.5$.
- [9] KEKB B Factory Design Report, KEK Report No. 95-1, 1995 (unpublished).
- [10] Belle Collaboration, K. Abe *et al.*, KEK Report No. 2000-4 [Nucl. Instrum. Methods Phys. Res. (to be published)].
- [11] Throughout this Letter, whenever a mode is quoted, the inclusion of the charge conjugate mode is implied.
- [12] Belle Collaboration, K. Abe *et al.*, "Measurements of Polarization and CP Asymmetry in $B \rightarrow J/\psi + K^*$ decays" (2001); BELLE-CONF-0105.
- [13] We use the QQ B meson decay event generator developed by the CLEO Collaboration (<http://www.lns.cornell.edu/public/CLEO/soft/QQ>) and GEANT3 for the detector simulation; CERN Program Library Long Writeup W5013, CERN, 1993.
- [14] Particle Data Group, D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- [15] The measured B lifetimes are $\tau_{B^0} = 1.547 \pm 0.021$ ps and $\tau_{B^+} = 1.641 \pm 0.033$ ps (statistical errors only).
- [16] θ_{tr} is defined as the angle between the ℓ^+ direction in the J/ψ rest frame and the z axis, where the x axis is defined as the direction of motion of the J/ψ in the $Y(4S)$ rest frame. The x - y plane is defined by the K^* decay products in the J/ψ rest frame.
- [17] BABAR Collaboration, B. Aubert *et al.*, preceding Letter, Phys. Rev. Lett. **87**, 091801 (2001).